



ESTABLISHED FLORIDA LAWN SPECIAL
17-2-17 FLORIKAN®
PROFESSIONAL LANDSCAPE FERTILIZER
Staged Nutrient Release (SNR®)



FEATURES

- + 80% POLYMER COATED CONTROLLED RELEASE
- + CONTAINS SULFUR BASED KICKER
- + LOW P CONTENT
- + HIGH MAGNESIUM CONTENT AND MICRONUTRIENTS
- + COVERS UP TO 5,000 SQUARE FEET

BENEFITS

- + LASTS A FULL 6 MONTHS IN FLORIDA CONDITIONS
- + QUICK GREEN-UP IN COASTAL SOILS
- + REDUCES POTENTIAL RUN-OFF
- + MAINTAINS HEALTHY COLOR
- + EASILY APPLIED ONLY TWICE PER YEAR

GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	17.00%
6.50% NITRATE NITROGEN	
5.50% AMMONIACAL NITROGEN	
5.00% WATER SOLUBLE NITROGEN	
PHOSPHATE (P2O5)	2.00%
SOLUBLE POTASH (K ₂ O)	17.00%
CHLORINE (Cl) NOT MORE THAN	1.00%
TOTAL COPPER (Cu)	.02%
TOTAL MAGNESIUM (Mg)	2.20%
1.50% WATER SOLUBLE MAGNESIUM (Mg)	
IRON (Fe)	.80%
0.10% CHELATED IRON	
0.11% WATER SOLUBLE IRON (Fe)	
MANGANESE (Mn)	0.37%
0.01% WATER SOLUBLE MANGANESE	
ZINC (Zn)	0.05%
0.01% WATER SOLUBLE ZINC (Zn)	

DERIVED FROM: Polymer Coated Ammonium Nitrate, Polymer Coated Urea, Polymer Coated Potassium Nitrate, Polymer Coated Sulfate of Potash Magnesia, Ammonium Sulfate, Ammonium Phosphate, Sulfate of Potash Magnesia, Copper Sulfate, Iron EDTA, Magnesium Sulfate, Iron Sulfate, Manganese Sulfate, Zinc Sulfate

15.00 Units = Slow Release Nitrogen
 16.00 Units = Slow Release Potassium
 1.00 Units = Slow Release Magnesium

17-2-17-180-DYN

CAUTION!

KEEP OUT OF REACH OF CHILDREN
 HARMFUL IF SWALLOWED
 This material contains Copper and Manganese Sulfate, breathing dust may be harmful to respiratory system, nervous system, and kidneys.

ROTARY SPREADER SETTINGS (LBS. /1000 SQ. FT.)

POUNDS OF FERTILIZER PER 1000 SQUARE FEET ANNUALLY

Low	Medium	High
10	11	12 1/2

DIRECTIONS FOR USE

FLORIKAN fertilizers are specially formulated for use on sports turf. Rotary spreader settings vary with different makes of equipment. Settings will be determined by testing a small area with the user's equipment. For Turf use a rotary spreader or chest spreader application method to apply the fertilizer. Apply one to two times a year as needed to maintain plant growth. You may need to adjust the setting depending on walking speed, type of spreader, conditions, and product.

MANUFACTURED FOR: ENVIRO-SAFE LABORATORIES, INC.
 1579 BARBER RD. SARASOTA, FL 34240,
 PHONE: 800.322.8666 FAX: 941.377.3633

F1130

NET WEIGHT
50 LBS.
(22.7 KG)

Blue Label
 Revised 5/22/07

CONTAINS:

